



Incidence of Urinary Tract Infections among Adolescent and Adult Women

Onwujekwe Emmanuel Chijioke^{1,*}, Ezemba ConstanceChinyere¹ and Agbo Angela²

¹Department of Microbiology, Coal City University, Enugu, Nigeria

²Microbiology Laboratory Department, Coal camp Health Centre, Enugu, Nigeria

Abstract

This study was to determine the bacteria that are prevalent in Urinary tract infections and the sensitivity of organisms isolated from the urine of adolescent and adult women of 8- 52 years old in Ogbete Coal camp in Enugu metropolis, south east Nigeria. A total of 218 individuals, both in and out patients were investigated, 197 (90.3%) women were diagnosed for Urinary tract infections. Therefore, incidence of UTI were prevalent in the age groups of 13-17 and 18-22, with P value (p=0.047 and S.D of ± 5.6). The microorganisms implicated in infection were bacteria, fungi, yeast and protozoa. Among the bacteria, two were identified as Gram positive cocci i.e. *Staphylococcus aureus* and *Staphylococcus saprophyticus*, while Gram negative bacilli were *Escherichia coli*, *Klebsiella*, *Proteus*, *Pseudomonas aureginosa*. Yeast like fungus *Candida albican*, protozoan (*Trichomonas vaginalis*) was also identified. The isolated organism which was prevalent was *Staphylococcus aureus* (48.2%); followed by *Escherichia coli* (34.5%). Also, *Pseudomonas aureginosa*, *Proteus* and *Klebsiella* were isolated. Factors responsible for frequent cases of Urinary tract infections among diagnosed women include shortness of urethra among the females, lack of personal hygiene, sexual intercourse, socio economic challenges in various homes and others. Sensitivity profiles revealed that Augumentin was highly susceptible to *Staphylococcus aureus* (65%) while *Escherichia coli* were highly resistant to commonly used drugs such as Ampicillin, and Cloxacillin while Gentamycin was the most active against bacteria isolates.

Keywords: Urinary Tract Infection, Adolescent Women, Antimicrobial Sensitivity, *Staphylococcus aureus*, *Escherichia coli*

Article Info:

Received:

June 2, 2018

Received Revised:

June 28, 2018

Accepted:

July 10, 2019

Available online:

July 25, 2019

*Corresponding author:

constancechinyere790@gmail.com

How to cite:

Chijioke OE, Chinyere EC, Angela A. Incidence of urinary tract infections among adolescent and adult woment. *Abasyn Journal of Life Sciences* 2019; 2(1): 34-41.

1. INTRODUCTION

Urinary tract infection (UTI) is an infection in any part of the urinary system – kidneys, ureters, bladder and urethra ¹. Worldwide, about 152 million people are diagnosed with urinary tract infections each year with morbidity of about 196,500 (WHO; 2017)². In developing countries like Nigeria, the rate of UTI's is high especially among women ³. Bacteriuria is defined as the presence of $>10^5$ colonies of a single pathogen per milliliter of urine. Urinary tract infection is more common in females than males as a result of opening of the urethra near the anus and shortness of urethra in females ⁴. According to Akobiet *al* ⁵, urine of females

has more suitable pH and osmotic pressure for the growth of *Escherichia coli* than urine from males. Urinary tract infection is classified into six categories as uncomplicated infection, complicated infection, isolated infection, unresolved infection, re-infection, and relapse infection⁶. Notable clinical symptoms of urinary tract infections include lower abdominal pain, urgency and frequency of micturition, dysuria, itching, pyuria, formation of blisters and ulcers in genital areas. Bacteria are the main causes of urinary tract infections among individual worldwide⁷. Urinary tract infections are commonly caused by bacteria such as *Escherichia coli*, *Klebsiellasp*, *Proteus*, *Enterococcusfaecalis*, and *Enterobacter*, *Pseudomonas aureuginosa*, *Staphylococcus aureus*. Group B *Streptococcus* is usually reported with increased rates in patients with urological disorders and following repetitive course of antibiotic treatment. The prevalence and the degree of occurrence of one or two of these organisms are dependent on the environment (Omonighoet al,¹). High sexual activities, poor hygiene, low economic status, are some risk factors that predispose adolescent and adult women with urinary tract infections⁹. According to Azubike et al,¹⁰ commonest mode of infection is the ascending route, through which organisms of the bowel flora contaminated the urethra, ascend to the bladder and migrate to the kidney or prostate. Knowledge about the type of pathogens responsible for Urinary tract infections and their susceptibility patterns may help clinicians to choose the right empirical treatment¹¹.

Therefore, this study aims to determine the bacteria that are prevalent in urinary tract infections in adolescent women and adult women, and also the antimicrobial susceptibility profiles of drugs used in the treatment of such infections.

2. MATERIALS AND METHODS

2.1 Study Area

Ogbete Coal camp area of Enugu is located in Enugu North local government in the heart of Enugu. The area is densely populated as a result of its proximity to Ogbete main market, which is the largest market in Enugu state. Also the presence of an automobile market at the Ogbete Coal camp town contributed to the dense population of this area. Therefore, the people of this area are mainly traders, artisans and students. There is government owned institutions as well as hotels around the area.

2.2 Study Population

The research was a retrospective studies carried out between May 2016 and May 2017 and was conducted in the microbiology section at Ogbete Coal camp Health Centre for urinary tract infections issues. This got the approval of the ethical committee of the Health Centre. Verbally informed consent was obtained from all patients prior to specimen collection and the permission to that effect was obtained from the ethical committee of the health center. Their urine sample was obtained through a mid-stream catch technique.

2.3 Sample Collection

Two hundred and eighteen (218) mid-stream urine (MSU) samples were collected inside sterile bijou bottles from the patients. They were instructed on how to collect the samples. The samples were labelled and processed within 30 minutes to 1 hour of sample collection.

Ten ml of each well mixed urine sample was centrifuged at 2000g for 5 minutes. The supernatant was discarded and a drop of the deposit was examined microscopically at high magnification for pus cells, red blood cells, epithelial cells, cast, crystal, yeast-like cells and *Trichomonas vaginalis*. Pus cells > 5 per high power field were significant in an individual.

2.4 List of Materials and Reagents

Materials used in the study include spatula, bijou bottles, stopper, masking tape, conical flask, autoclave, gas cylinder, bunsen burner, incubator, refrigerator, electronic weighing balance, colony counter, marker, Petri dish, wire loops, slides, test tube glass, pipette, centrifuge, microscope. Nutrient agar, MacConkey agar, blood agar, distilled water, crystal violet, lugol iodine, acetone, caramel, and antimicrobial sensitivity disc.

2.5 Sample Processing

Urine sample were cultured on MacConkey agar and Blood agar. The agar were prepared and sterilized based on the manufacturer's instruction using the autoclave. After sterilizing, the media was allowed to solidify at a temperature of 50⁰C-55⁰C, then it was dispensed into sterile Petri dish. The inoculation was carried out and the media incubated for about 24hours at 37⁰C. The organism grown was sub-cultured so as to isolate pure cultures of organisms. A count of >.10⁵CFU (Colony forming unit) was considered significant to indicate UTI. Isolated bacterial species were characterized by Gram staining followed by microscopic examination, motility and biochemical tests. Bacteria identification was based on standard culture and biochemical characteristics of the isolates as described by Omer and Fadil¹² and Cheesbrough⁴.

2.6 Antimicrobial Sensitivity Testing

The susceptibility test was carried out using disc diffusion technique as recommended by National Committee for Clinical Laboratory Standards (NCCLS)¹³. The bacterial isolates and standard strains of *Escherichia coli* ATCC 25922, *Enterococcus faecalis* 29212, *Pseudomonas aureuginosa* ATCC 27853 were used for quality control. The antimicrobials used are: Ceftriaxone, Ampicillin, Cloxacillin, Augumentin, Streptomycin, Azythromycin, Gentamycin, Cefuroxime, Cotrimoxazole, Nitrofurantoin, Nalidixic Acid.

Statistical Analysis: This data was analysed using SPSS version 20.0.

3. RESULTS AND DISCUSSIONS

A total of 197 (90.3%) out 218 samples collected were confirmed positive with urinary infections. The age distribution of urinary tract infections in women are shown in Table 1. Out of the nine age groups represented in the table, positive bacterial isolates were prevalent in individuals between 18-22 and 13-17 years. Bacterial isolates were least in individuals between the age group of 47-52. The bacterial species implicated in the urinary tract infections are *Escherichia coli*, *Klebsiella sp*, *Staphylococcus saprophyticus*, *Staphylococcus aureus*, *Proteus mirabilis* and *Pseudomonas aureuginosa*. The frequency of occurrence of isolates is presented on Table 2. *Staphylococcus aureus* was found to be the most prevalent organism among the isolates, with 48.2% occurrence. This was followed by *Escherichia coli* which accounted for 34.5 % of the isolates. *Staphylococcus saprophyticus* was least frequently isolated.

Table 3 showed varying antimicrobial susceptibility pattern of isolates to the various antimicrobials used. Gentamycin was highly effective against most bacterial isolates. Ampicillin was virtually useless against all bacterial isolate. Table 5 showed the characteristics of bacteria isolates. Table 6, showed the microscopic findings of urine deposits.

Mid-stream catch technique was used to ensure those normal floras were flushed before the samples for analysis were obtained⁴. Frequent voiding is necessary to flush out invading bacteria from the walls of the urethra^{4,14}.

The result of this study showed that among the population of 218 samples, 197 (90.3%) were diagnosed positive of bacterial infections. Therefore, incidence of UTI were prevalent in the age groups of 13-17 and 18-22, with P value (p=0.047 and S.D of ± 5.6) statistically not significant. The frequency of occurrence of isolates is presented on (Table 2). *Staphylococcus aureus* is a predominant pathogen which accounted for 48.2 % of the bacteria isolates that caused urinary tract infection in adolescent women; this may be attributed to high sexual activities among their age group [9]. *Escherichia coli* also had a significant number which accounted for 34.5% of the isolates that caused urinary tract infection in women. This is in line with the work of Doud *et.al*¹⁵. The reason for the high prevalence of *Escherichia coli* can also be attributed to the suitable pH and osmotic pressure provided by the female urine⁵. *Klebsiella sp* was responsible for 6.6% of the isolates. *Proteus* species accounted for 4.0%; a Gram- negative bacterium and flora of the intestinal tract that causes infection when it makes its way from the intestine to urinary bladder and urethra. *Staphylococcus saprophyticus* was the least among the bacteria isolates with 1.5% occurrence. However, this result agrees with the findings of MMDT, that about 40% of the urinary tract infections are caused by Gram negative species¹⁶.

This study (Table 1) reports high incidence of UTIs between age groups 18 and 22 years (98.1%), and 13-17 years (96.1 %). The most common uropathogen isolated in this study was *Staphylococcus aureus*. This findings is similar to two reports in Benin City (urban settlement)^{17, 18}, and findings of Akinola *et al*¹⁹. These reports were on asymptomatic bacteriuria against the present study that deals on symptomatic bacteriuria.

However, our findings are contrary to the findings of Bankole *et al*⁹, and Akobi *et al*⁵, who reported *Escherichia coli* as the predominant isolates.

The high incidence of urinary tract infection in women as earlier mentioned is as a result of shortness of urethra, high sexual activity and closeness of urethral opening to the anus⁹. Also, socio-economic factors, is also a major risk factor as mentioned earlier¹⁰. Some homes where some of the studied population lived, were highly polluted environment, with unsuitable toilet facilities thereby exposing such inhabitants to numerous pathogenic microorganisms.

Antibiotic abuse and practicing incomplete antibiotic dosage has highly promoted the dissemination of multi drug resistant bacteria²⁰. *Staphylococcus aureus* was the highest uropathogen, showed high susceptibility to Nitrofurantoin and least susceptible to Cloxacillin. *Staphylococcus saprophyticus* followed similar pattern of susceptibility with *Staphylococcus aureus*. The present study found the *Escherichia coli* and *klebsiella* species isolates to be highly resistant to the commonly used drugs such of the Penicillin group such as Ampicillin and Cloxacillin. Ahmed *et al*⁶ reported that Amoxicillin, Ampicillin, and Sulfanomides are no longer the drug of choice for empirical treatment because of widespread emergence of resistance in 15-20% of *E.coli* in several areas of USA and other countries. However, the antimicrobial susceptibility profile indicates that Nitrofurantoin (69.1%) were the most active agents against *Escherichia coli*. Aminoglycosides, example (Gentamycin and Augumentin) are the next drugs that are most susceptible to the isolates by (58.8%) and (51.4%) respectively. This result is in agreement to the recommendations of Akobi *et al*,⁵. *Klebsiella pneumonia* was susceptible to Nitrofurantoin and Cefuroxime both by (76.6%), but was completely resistant to Ampicillin. *Proteus* was highly susceptible to Gentamycin and Ceftriaxone but was also highly resistant to Streptomycin, Ampicillin, and Nitrofurantoin. *Pseudomonas aureuginosa* was moderately susceptible to Gentamycin and Ceftriaxone. *Proteus* sp and *P.aureuginosa* isolates were resistant to most antimicrobials. Most bacterial isolates were least susceptible to Cotrimoxazole. This study is similar to the findings by Mezue *et al*,²¹ at University of Nigeria found Cotrimoxazole to be virtually useless against uropathogen. The resistance of most of the isolates to the penicillin group could be as a result of long term use of these drugs over the years.

However, patients should be advised and encouraged to drink plenty of fluids (two to three liters per day) and to urinate frequently to help flush bacteria from the bladder. Avoiding multiple sexual partners will reduce the risk of both UTIs and sexually transmitted infections. Women are encouraged to avoid spermicidal contraceptive, diaphragms and vaginal douching, which may irritate the vagina and urethra; facilitating the entry and colonization of bacteria within the urinary tract. Meanwhile, some recent studies by Ahmed *et al*^{6, 22, 23, 24, 25} have suggested alternative therapies other than antimicrobials in the treatment of RUTI'S. Some of the measures include the use of Cranberry juice and tablets, acupuncture, probiotics such as *Lactobacillus* found in fermented milk products; mainly yoghurt, as well as immunoprophylaxis. Nevertheless, more clinical studies need to be carried out to determine their role in RUTI prevention.

Table 1. Incidence of UTI's in relation to Age among adolescent and adult women

Age interval	No Tested	% positive	% negative
8-12	19	11 (57.9)	8 (42.1)
13-17	51	49 (96.1)	2 (3.9)
18-22	53	52 (98.1)	1 (1.9)
23-27	11	8 (72.7)	7 (63.6)
28-32	37	35 (94.6)	2 (5.4)
33-37	14	10 (71.4)	4 (28.6)
38-42	8	6 (75)	2 (25)
43-47	14	11 (78.6)	3 (21.4)
48-52	10	5 (50)	5 (50)

Table 1; showed prevalence of infections was most among women between the ages of 18-22 and 13 and 17 (98.1% and 96.1%) positive cases respectively. Age group 48-52, had the least incidence rate with 5 (50%) positive cases.

Table 2. Prevalence of organisms isolated from the urine

Organism	Number of patients	Number of occurrence (%)
<i>Escherichia coli</i>	197	68 (34.5%)
<i>Klebsiella</i>	197	13 (6.6%)
<i>Staphylococcus saprophyticus</i>	197	3 (1.5%)
<i>Staphylococcus aureus</i>	197	95 (48.2%)
<i>Proteus</i>	197	8 (4.0%)
<i>P. aureuginosa</i>	197	11 (5.6%)

Staphylococcus aureus was most prevalent in the urine samples (48%), next was *Escherichia coli* 34% and the least prevalent was *Staphylococcus saprophyticus*.

Table 3. Susceptibility Profiles of Urinary Bacterial Isolates

ANTIBIOTICS	<i>E.coli</i> (N=68)	<i>K. pneumonia</i> (N=13)	<i>S.saprophyticus</i> (N=3)	<i>S.aureus</i> (N=95)	<i>P.mirabilis</i> (N=8)	<i>P.aureuginosa</i> (N=11)
CEFT	26 (38.2%)	4 (30.7%)	1 (33.3%)	19 (20%)	6 (75%)	0(0.0%)
AMP	2 (2.9%)	0 (0.0%)	0 (0.0%)	4 (4.2%)	0 (0.0%)	NA
CLOX	NA	NA	1 (33.3%)	14 (14.7)	NA	NA
AUG	35 (51.4%)	4 (30.7%)	2 (66.6%)	62 (65.2%)	0 (0.0%)	0 (0.0%)
STREPT	20 (29.4%)	6 (46.1%)	1 (33.3%)	20 (21.0%)	0 (0.0 %)	4 (36.3%)
AZY	29 (42.6%)	4 (30.7%)	1 (33.3%)	32 (33.6%)	6 (75%)	0 (0.0%)
GENT	40 (58.8%)	4 (30.7%)	1 (33.3%)	29 (30.5%)	6 (75%)	4 (36.3%)
CEF	7 (10.2%)	10 (76.9%)	0 (0.0%)	18 (18.9%)	0 (0.0%)	0 (0.0%)
COT	9 (13%)	2 (15.3%)	1 (33.3%)	19 (20%)	4 (50%)	0 (0.0%)
NIT	47 (69.1%)	10 (76.9%)	2 (66.6%)	47 (49.4%)	0 (0.0%)	NA
NAL	27 (39.7%)	4 (30.7%)	0 (0.0%)	16 (16.8%)	0 (0.0%)	NA

Bacteria Isolates with Zone of inhibition = 16mm is sensitive while < 15mm is resistant to a particular antibiotics. CEFT= CEFTRIAZONE. AMP= AMPICILLIN. CLOX= CLOXACILLIN. AUG= AUGUMENTIN. STREPT=STREPTOMYCIN. AZY= AZYTHROMYCIN. GENT= GENTAMYCIN. CEF=CEFUROXIME. COT= COTRIMOXAZOLE. NIT= NITROFURANTOIN. NAL=NALIDIXIC ACID. N= TOTAL NUMBER.

Table 3; showed varying antimicrobial susceptibility pattern of isolates to the antimicrobials used.

A total of 42 samples were resistant to either one or two antimicrobials used in bacteria susceptibility testing. *Staphylococcus aureus*, the most frequently isolated bacteriuria showed high susceptibility to Augumentin (65.2%) and Nitrofurantoin (49.4%) and least susceptible to Ampicillin (4.2%). *Escherichia coli*, second frequently isolated bacteriuria was susceptible to Nitrofurantoin (69.1%) and Gentamycin (58.8%) but least susceptible to Ampicillin (2.9%). *Klebsiella sp* was susceptible to Ceftriaxone and Nitrofurantoin both by (76.9%), however was completely resistant to ampicillin by (100%). *Staphylococcus saprophyticus* was susceptible to Augumentin and Nitrofurantoin both by (66.6%) and completely resistant to Ampicillin and Nalidixic acid both by (100%). *Proteus mirabilis* was susceptible Ceftriaxone, Azythromycin, and Gentamycin by (75.5%) but showed (100%) resistance to Ceftriaxone, Augumentin, Streptomycin, Cefuroxime, Nitrofurantoin and Nalidixic acid. *Pseudomonas aureuginosa* was moderately susceptible to Streptomycin and Gentamycin both by (36.6%) and resistant to other antimicrobials.

Table 4. Pattern of bacterial occurrence in age groups

Age group	<i>E.coli</i>	<i>klebsiella</i>	<i>S.aureus</i>	<i>proteus</i>	<i>p.aureuginosa</i>	<i>S.saprophyticus</i>
8-12	3	1	8	-	-	-
13-17	14	1	33	-	-	1
18-22	20	5	22	2	1	2
23-27	3	1	4	-	-	-
28-32	16	2	11	2	6	-
33-37	7	-	2	-	1	-
38-42	1	2	3	4	1	-
43-47	4	1	6	-	2	-
48-52	2	-	4	8	-	-

Staphylococcus aureus was found to be main urinary pathogen among children between 8-11 years. It was observed that *Escherichia coli* and *Staphylococcus aureus* were implicated in infections in adolescents between 18-22 years as well as in women between 43-47, and 48-52.

Table 5. Identification of characteristics of bacterial isolates

Catalase	Coagulase	Oxidase	Urease	Indole	Glucose	Sucrose	Lactose	Organism
+	-	-	-	+	A	A	AG	<i>Escherichia coli</i>
+	ND	-	+	-	AG	A	AG	<i>Klebsiella sp.</i>
+	+	-	-	ND	A	A	A	<i>S.aureus</i>
-	ND	-	+	+	-	A	ND	<i>Proteus</i>
ND	ND	+	-	ND	ND	ND	ND	<i>P.aureuginosa</i>
+	-	-	-	ND	A	A	A	<i>S.saprophyticus</i>

Key: +=Positive, -=Negative, A=Acid, G=Gas, ND= Not Done.

Table 6. Microscopy of urine deposits

Age	Appearance	WBC	RBC	Yeast	Epithelia cells	<i>T.vaginalis</i>
8-12	Amber, & cloudy	11-20	F	F	F, M	F
13-17	Amber, red & cloudy,	11-40	MD	F	M	F
18-22	Amber, red & cloudy	11-40, 40-	F, M	M	F	MD
23-27	Amber, red & cloudy	11-40	F, MD, M	M	M	MD
28-32	Amber, red & cloudy	11-40, 40-	MD, M	M	MD	F
33-37	Red & cloudy	11-40, 40-	F, M, MD	M	F	M
38-42	Red & cloudy	11-40, 40-	MD	M	M	MD, M
43-47	Amber red & cloudy	11-40, 40-	M	M	M	MD
48-52	Amber, red & cloudy	11-40	M	M	F	MD

Key: F = Few, MD = Moderate, M = Many

In urine microscopy, white blood cells (pus cells) were found. Up to 10WBCs / HPF (high power field, i.e. using 40x objectives) was considered as few in number. Moderate number: 11-40/ HPF, and more than 40 WBC/ HPF, was considered as many.

4. CONCLUSIONS

Staphylococcus aureus is a predominant organism causing urinary tract infections in adolescent and adult women as a result of unprotected sex and high sexual activity at their age. *Escherichia coli* are also frequently isolated organism that causes UTIs. Urinary tract infections occur mostly in females than their male counterparts. It has dangerous effect when left untreated. The prevalent isolates were susceptible to Augmentin, Nitrofurantoin and Gentamycin. This study advocates early diagnosis of urinary tract infection, regular monitoring of drug resistant phenotype of UTI pathogens to improve public health treatment and reduce cases of infections with other complications caused as a result of urinary pathogens in our society.

ACKNOWLEDGEMENTS

We are very grateful to the entire staff of the Faculty of Natural and Applied Science Coal City University Enugu, and Ogbete Coal camp Health Centre Enugu for their cooperation and for given us access to their research laboratories.


CONFLICT OF INTEREST

All authors declare no conflict of interest regarding this article.

REFERENCES

1. Omonigho, S.E., Obasi, E.E. & Akukalia. R.N. In Vitro Resistance of Urinary Isolates of *Escherichia coli* and *Klebsiella* species to Nalidixic Acid, 2001
2. World health organization (WHO). Accessed on 24th June, 2017.
3. Orret, F. A. & Davis, G.K. A comparison of the antimicrobial susceptibility profile of urinary pathogens for the years 1999 and 2003. West Indian Medical Journal, 2006 55: 95-99.
4. Cheesbrough. M. District Laboratory Practice in Tropical Countries part 11, Cambridge University press, UK, 2004.
5. Akobi, O.A., . Inyinbor, H.E., Emumwem, E.G, Ogedengbe, S O., Uzoigwe, E.O, Abayomi, R.O., Emumwem, E.F., OkorieEbie, M.Y., Kandakai-Olukemi,J. Y. T. & Ayanbadejo, K.B. Tanyiga. Urinary tract infections in a Nigeria military Hospital Niger. Journal of Microbiology., 2001; 15(1): 31-37.
6. Ahmed, A.& Ghadeer,V. Recurrent Urinary Tract Infection Management in Women, Sultan Qaboos University Medical Journal, 2013.
7. Ebie, M.Y., Kandakai-Olukemi, Y. T., Ayanbadejo, J., & Tanyiga K.B. Urinary tract infections in a Nigeria military Hospital, Nigerian Journal of Microbiology., 2001;15(1): 31-37.
8. Guentzel, M.N. *Escherichia klebsiella, Enterobacter, Serratia, Citrobacter, and Proteus*. Barron's Medical Microbiology 4th ed. Univ of Texas Medical Branch, 1996.
9. Oladeinde, B H., Omoregie, R. Olley, M. & Anunibe, J.A. Urinary tract infection in a rural community in Nigeria. North American Journal of Medical Sciences, 2011; 3 (2): 75-77.
10. Azubike, J.C., & Nkeaniginieme, K.E.O. Paediatrics and Applied Health in Nigeria, 1999.
11. Ogbukagu, C.M., Anakwenze, V.N., Ekwealor, C.C., Ezemba, C.C. & Ekwealor, I.A. Incidence of Urinary tract Infections (UTI) among patients attending Primary Health Centers in Anambra State, 2016.
12. Omer, E.I. & Fadil. E. Principles of Medical Microbiology. University students Library, Makkah AL Mukarramah edition, 1986.
13. NCCLS, author. Performance Standards for Antimicrobial - Susceptibility Testing, 11 th informational Supplement, NCCLS document M100-S11. Wayne, Pennslyvia: National Committee for Clinical Laboratory Standards. (NCCLS); 2001.
14. Bankole, O.H, Richard, O., Mitsan, O.& Joshua, A.A. Urinary tract infection in a rural community in Nigeria. North

- American Journal of Medical Sciences,2011; 3 (2) 75-77.
15. Doudz, Z. & Affif.,C. *Escherichia coli* Isolated from Urinary Tract Infections of Lebanese Patients between 2000 and 2009, Epidemiology and Profile of Resistance, 2011.
 16. MMDT; Bacterial Infections, (2005); Merk Manual.
 17. Omoregie, R. , Erebor, J.O., Ahonkai, I., Isobor, J.O. & Ogefero H.O. Observed changes in the prevalence of Uropathogen in Benin City, Nigeria. NZJ Med Lab Sci.,2008; 62: 29-31.
 18. Omoregie, R. & Eghefona, N.O. Urinary tract infection among asymptomatic HIV patients in Benin City, Nigeria. Br. J Biomed Sc., 2009; 66(4): 190-193.
 19. Akinola, B.A., Nwabuisi, C, Aboyeji, A.P., Ajayi, N.S., Adeola, F.& Olurotimi, O.F. Asymptomatic Bacteriuria in Antenatal Patients in Ilorin, Nigeria. Oman Medical Journal,2012 ; 27(1):31-35.
 20. Rahman, S R., Ahmed, M.F. & Begum,V. Occurrence of Urinary Tract Infection in Adolescent and Adult Women of Shanty town in Dhaka City, Bangladesh.2014; 24 (2) 145-154.
 21. Mezue, K., Ofong, C. Nmezi, D., Ugochukwu-Obi, G G. Antibiotic Sensitivity Patterns in Urinary Tract Infections at a Tertiary Hospital. Journal of the University of Nigeria Medical Students, 2006.
 22. Schmidt, D.R & Sobota, A.E. An examination of the antiadherence activity of Cranberry juice on urinary and non-urinary bacterial isolates.1988; 55:173-181, 1988.
 23. Alraek, T., Soedal, L.I.F.,Fagerheim, S.U., Digranes, A.,& Baerheim A. Acupuncture treatment in the prevention of uncomplicated recurrent lower urinary tract infections in adult women. American Journal Public health., 2002 92: 1609-11.
 24. Reid, G. & Bruce, A.W.. Probiotics to prevent Urinary tract infections: The rationale and evidence. World Journal Urology.,2006; 24: 28-32.
 25. Kul', E.V. & Chavenia, A.A. Efficacy of Uro-vaxom in recurrent infectious inflammatory disease of the urogenital



This work is licensed under a Creative Commons Attribution-Non Commercial 4.0 International License. To read the copy of this license please visit:

<https://creativecommons.org/licenses/by-nc/4.0/>