



A New Record of Genus *Heteropternis* Stal 1873 (Orthoptera: Acrididae: Oedipodinae) from Pakistan

Waheed Ali Panhwar^{1*} and Shoaib Ali²

¹Department of Zoology, Shah Abdul Latif University Khairpur, Mir's Sindh, Pakistan

²Department of Zoology, Hazara University, Mansehra, Pakistan

Abstract

Grasshoppers belonging to family Acrididae are vital herbivores that can severely disturb the growth of crops, their nutritional values and production rate. Band-winged grasshoppers belonging to genus *Heteropternis* Stal, 1873 were collected from different localities of Hazara Division during the year of 2016. *Heteropternis respondens* (Walker, 1859) was recorded for the first time from Pakistan. Additionally, photograph along with line drawing is provided for the first time. Finding of present study will contribute to the biodiversity of Pakistan.

Keywords: *Heteropternis resprodens*, New record, Identification, Band-winged, Distribution, Pakistan

Article Info:

Received:

May 17, 2018

Received Revised:

May 29, 2018

Accepted:

May 29, 2018

Available online:

May 30, 2018

***Corresponding author:**

waheed.panhwar@salu.edu.pk

How to cite:

Panhwar WA, Ali S. A New Record of Genus *Heteropternis* Stal 1873 (Orthoptera: Acrididae: Oedipodinae) from Pakistan. *Abasyn Journal of Life Sciences* 2018; 1(1): 36-41.

1. INTRODUCTION

Grasshoppers belonging to family Acrididae are important herbivores that can intensely disturb the growth of crops¹, their nutritional values and production rate². The subfamily Oedipodinae includes grasshoppers generally known as "winged band grasshoppers"³. They are generally found in high meadows, bushes, open forest, agronomic lands and fields⁴. There are 135 reported genera with 285 described species of family Acrididae and 23 genera with 52 species of Oedipodinae from India⁵. Moreover 5 species were recorded from Epacromiimi relating to the genera *Aiolopus* Fieber, 1853, *Heteropternis* Stål, 1873 and *Hilethera* Uvarov, 1923⁴. The genus *Heteropternis* Stål, 1873 includes 21 species worldwide out of which 10 are reported from Asia⁶ and *Heteropternis respondens* (Walker, 1859) is the only species which is reported from India⁷.


Periodic faunastical studies on insect fauna had been made from different regions of Pakistan some of which are reported by⁸⁻²⁷ however, they did not record *Heteropternis* Stal, 1873 from this region.

Present study will be the contribution as a first record of genus *Heteropternis* from Pakistan. Which is based on external morphological studies and evaluation of previous literature with description of various species of *Heteropternis*.

2. MATERIALS AND METHODS

2.1 Location

Specimens were collected from different localities of Hazara division. It is located at 34°30'0" North and 73°15'0" East. Hazara division is bounded by northern areas at north; Azad Kashmir is in its east, Islamabad is situated at the south of Hazara division. Total area of Hazara division is 18,013 km². It is the wettest part of Pakistan. Hazara division has an unusual bimodal rainfall regime with one peak in February and another in July and August. As Hazara is present at high altitude, temperature in Hazara division is cooler than on plains (Map.1).



Map.1. Showing surveyed areas

2.2 Sampling, Killing and Preservation

Samples were collected from August 2016 to March 2017. Specimens were killed using Potassium cyanide into glass jars. The insects were then dried and kept into insect boxes by keeping naphthalene balls in order to prevent from parasitoids and predators.

2.3 Identification

Identification of specimens was carried out using stereoscopic dissecting binocular microscope with the help of keys and description available in literature and on website <http://orthoptera.speciesfile.org/>

2.4 Depository

The material is deposited at Insect Museum, Department of Zoology, Shah Abdul Latif University and Hazara University, Mansehra.

2.5 Material examined

Several specimens from Hazara Division.

3. RESULTS AND DISCUSSIONS

3.1 Taxonomy

Family Acrididae MacLeay, 1821

Subfamily Oedipodinae Walker, 1871

Tribe Epacromiini Brunner von Wattenwyl, 1893

Genus *Heteropternis* Stål, 1873

Type species: *Acrydium respondens* Walker, 1859

3.2 Genus Diagnosis

Size medium, from slender. Antennae filiform, slightly thickened towards the apex, reaching the hind margin of pronotum or somewhat longer. Head rugulose punctate. Frontal ridge convex in profile and slightly reclinate, distinctly narrowed at the fastigium, gradually widened towards clypeus; its surface rugose foveolate, impressed above and a ocellum. Fastigium of vertex sloping and forming a somewhat angulated arch with the frontal ridge, broader than long, narrowed in front, its surface concave, without media keel, separated from the vertex by an incomplete keel or rugosity; foveolae temporales small, indistinct, acutely triangular. Lateral facial carinae angulately rounded. Pronotum short, constricted in the prozona, rugose punctured, median keel well developed, sharp; transverse sulci indistinct in prozona, third sulcus distinct, though very fine, both on the disc and on the lobes, cutting the median keel before the middle; lateral keel absent or only perceptible in the metazoan as a short oblique rugosity; front angle obtuse, hind angle obtusely angulated with the sides straight; lateral lobes distinctly shorter than high, rugose punctured, especially in the hinder part, front margin slightly rounded, front angle obtuse, lower margin straight and oblique backwards to beyond the middle, hind angle more or less rectangular or rounded. Sternum with scattered punctures; mesosternal lobes transverse, their interspace as broad as or broader than one of the lobes; metasternal interspace transverse. Elytra extending beyond the hind knees, subcoriaceous in the basal third, hyaline in the rest, stigmatic arch about in the middle; costal area feebly expanded basally gradually narrowing apical and reaching the middle of elytra; scapular area reaching the apical third, coriaceous basally with transverse veinlets in the apical half; discoidal area densely venulated with transverse veinlets, except at the base where they are more irregular, intercalate vein well developed, anterior discoidal area distinctly larger than the posterior one; interlunar area narrower than discoidal, irregular venulated or with sparsely transverse veinlets, with more or less distinct false vein; all areas of the apical half of elytra with transverse veinlets forming elongate cellules. Wings hyaline, coloured basally, without dark transverse band. Hind femora robust, with the upper keel scarcely denticulate. Hind tibiae nearly straight, armed with 10 inner spines and 9 outer spines, without outer apical spine; the inner apical spurs longer than the outer ones, especially the lower inner which is about twice as long as the outer one; all spurs are hooked at their apex, but the lower inner is more abruptly hooked at apex. Hind tarsi short, the first joint subequal to the two other together. Supra-anal-plate of the male triangular with the sides slightly rounded and the apex obtuse; in the basal half a shallow median sulcus. Cerci cylindrical, apex obtuse and slightly curved, slightly longer than the supra-anal-plate. Subgenital plate short, obtusely conical. Supra-anal-plate of female triangular, narrow, apex subacute, with a shallow impression in the basal half. Cerci short, obtusely conical. Valves of ovipositor short, obtuse. Sub-genital-plate longer as broad, posterior margin rounded.

3.3 *Heteropternis respondens* Walker 1859

3.4 Description of species

General coloration brown, with reddish brown and yellowish marking. Antennae filiform, slightly thickened towards the apex, reaching the hind margin of pronotum or somewhat longer. Antennae yellowish brown, darker apically. Head brown with irregular blackish-spots, dots or stripes. Pronotum brown or greenish brown or reddish, with pale reddish brown dots on each side of the disc, or metazoan and posterior part lateral lobes, reddish or yellowish-brown or greenish, very variable. The lateral lobes of pronotum in the anterior part with two shining dots. Elytra brown with a pale, more or less triangular spot in the basal third and some little spots more; apical part tessellate with hyaline. Wings reddish or yellowish basally with the apex slightly infumated. Hind femora yellowish brown on their outer face with irregular rows of dark spots along the keels; with 2-3 irregular black spots above; inner face and lower sulcus reddish, sometimes with dark fasciae or spots. A small preapical projection at the inner spur of the inner side of the hind tibia. Knees with brown and black variegated. Sternum and abdomen yellowish, margins of lobes and sternites darker. Hind tibiae red, with a small yellow ring basally, the extreme tip infuscated, spines reddish with black tips, spurs yellowish-brown with black tips. Cerci short, obtusely conical. Valves of ovipositor short, obtuse. Sub-genital-plate longer as broad, posterior margin rounded. (Fig1 a-f).

3.5 Morphometry (in mm)

♂: Total body length: 18-19; Length of pronotum: 4-4.5; Length of tegmina: 18-19; Length of femur: 11-12, ♀: Total body length: 21-23; Length of pronotum: 5; Length of tegmina: 23-24; Length of femur: 13-14.




Fig. 1. *Heteropternis respondens* Walker 1859, a, c) Pronotum Lateral view, b, d) Same but Dorsal view, e, f) Habitus Lateral and Dorsal view

3.6 Remarks

Finding of *Heteropternis respondens* (Walker, 1859) constructed as a new record for Pakistan.

3.7 Distribution

This species is distributed in Bangladesh, China, Indonesia, Japan, Java, Malacca, Malaysia, Myanmar, Nepal, Philippines, Sri Lanka, Sumatra, Taiwan, Australia, India and Pakistan.

2. CONCLUSIONS

It is concluded from the present study that Pakistan is more diverse region to be explored for the biodiversity of *Heteropternis* species. Further survey should be carried out to find new records or new species of this genus from this area.

ACKNOWLEDGEMENTS

The authors are highly thankful to Mehwish Kainat, Izat Naz, Halimullah and all research fellows for their help in sampling and photography of samples.

CONFLICT OF INTEREST

All authors declare no conflict of interest regarding this article.

REFERENCES

1. Parihar DR. *Grasshopper pests of grazingland vegetation and their management in Indian Desert*. Central Arid Zone Research Institute; Jodhpur.1987
2. Soomro F, Sultana R, Wagan M, Abbasi A. Fauna of Acrididae (Orthoptera: Acridoidea) from Nara Desert District Khairpur. *Sindh University Research Journal-SURJ (Science Series)*,2015; 47(4).23-24
3. ElEla SA, ElSayed W, Nakamura K. Incidence of orthopteran species (Insecta: Orthoptera) among different sampling sites within Satoyama area, Japan. *Journal of Threatened Taxa*, 2012;4(3), 2476-2480.
4. Gupta SK, Chandra, K.A new species of the short-horned grasshopper genus *Heteropternis* Stål, 1873 (Orthoptera: Acrididae) from Central India. In *Annales de la Société entomologique de France (NS)* ,2016; 52 (4), 196-203
5. Shishodia MS, Chandra K, Gupta SK. An annotated checklist of Orthoptera (Insecta) from India. Zoological Survey of India.2010.
6. Cigliano MM, Eades DC, Otte D, Braun H. Orthoptera species file. Version 5.0/5.0. [cited 2018 Apr 2]. Available from: <http://orthoptera.species.file.org>
7. Kirby WF. The fauna of British India, including Ceylon and Burma, I. Orthoptera (Acridiidae). London: Taylor and Francis;1914; ix + 276 pp.
8. Bughio BA, Sultana R, Wagan, MS.A new species of the genus *Hilethera* Uvarov (Oedipodinae: Acrididae: Orthoptera) from Pakistan. *J.Cell Anim. Biol.* 2012;6: 29-32
9. Cardinale BJ, Diane S, Srivastava JE, Duffy PW, Justin D, Amy LM, Sankaran Jouseau C. Effects of biodiversity on the functioning of trophic groups and ecosystems. *Nature*, 2006;443: 989-992.
- 10.Panhwar WA, Riffat S, Wagan MS, Khatari I, Santosh K. Systematic study on the various Tribes of Phaneropterinae (Tettigoniioidea: Orthoptera) occurring in Pakistan. *Pakistan Journal of Zoology*, 2014; 46: 203-213.
- 11.Panhwar WA, Riffat S, Wagan MS, Santosh K. Notes on the distribution and morphological description of *Glyphonotus sinensis* Uvarov, 1939 (Orthoptera: Tettigoniinae: Glyphonotini) from Pakistan. *International Journal of Advanced Research*. 2013;1 (9): 679-682

12. Panhwar WA, Riffat S, Wagan MS, Santosh K. On the distribution and taxonomy of *Conocephalus* species (Orthoptera: Tettigoniodea: Conocephalinae) from Pakistan. *Journal of Biodiversity and Environmental Sciences*. 2013;3 (11): 171-176
13. Panhwar WA, Riffat S, Wagan MS, Wagan YS, Santosh K, Solangi FH. Taxonomy and Ecology of Genus *Euconocephalus* Karny, 1907 (Orthoptera: Tettigoniodea: Conocephalinae) from Pakistan. *International Journal of Advanced Research* .2014;2 (2)268-277
14. Panhwar WA. Studies on the Systematic and Ecological Status of Tettigoniodea (Ensifera) Of Pakistan. PhD Thesis Submitted To Department of Zoology, University of Sindh ,2015;1-242.
15. Chapman RF. The structure and wear of mandibles in some African grasshoppers. *Proc. zool. Soc. Lond.* 1964;142: 107-121.
16. Dadd RH. Feeding behavior and nutrition in grasshoppers and locusts. *Adv. Insect Physiol* 1963;1: 47-109
17. Isley FB. Correlation between mandibular morphology and food specificity in grasshoppers. *Ann entomol. Soc. Am Florida University, Florida*; 1944;1- 5.
18. Joern A, Lawlor LR. Food and microhabitat utilization by grasshoppers from arid grasslands: comparisons with neutral models. *Ecology*. 1980;61: 591-599.
19. Joern A. Feeding patterns in grasshoppers (Orthoptera: Acrididae): Factor influencing diet specialization. *Ecologia*; 1979;38:325-347.
20. Lomer CJ, Bateman RP, Johnson DL, Langewald J, Thomas M. Biological control of Locusts and grasshoppers. *Annu. Rev. Ent*; 2001;46:667-702.
21. Mahmood K, Idris AB, Salmah Y. Tetrigidae (Orthoptera: Tetrigoidea) from Malaysia with the description of six new species. *Acta Entomologica Sinica*. 2007;50: 1272-1284.
22. Nadia A, Irfan A, Tanveer H, Ishtiaq A. *American-Eurasian J. Agric. & Environ. Sci.* 2015; 15 (8): 1693-1699.
23. Otte D. The North American grasshopper. Vol. I: Acrididae: Gomphocerinae and Acridinae. Harvard University Press, Cambridge, MA, 1981;1-275.
24. Riffat S, Wagan MS. Super flood destroyed the natural habitat of many animals in Sindh province *Proc. 2nd International Conference of (IC Biour-life) Shah Latif Uni.* 2010; 277-279.
25. Riffat S, Wagan MS. A revision of the Genus *Heiroglyphus* Krauss 1877 (Hemiacridinae: Acrididae: Orthoptera) with description of one new specie from Pakistan. *Pakistan Journal of Zoology*, 2012;1: 43-50.
26. Riffat S, Wagan YS, Wagan MS. Orthopteran Biodiversity of Thar desert, Sindh, Pakistan. *Pakistan Journal of Zoology*, 2013; 2: 299-304.
27. Uvarov BP. XI.—A revision of the Old World Cyrtacanthacrini (Orthoptera, Acrididæ).—I. Introduction and key to genera. *Journal of Natural History*, 1923; 11(61), 130-144.